

WHAT TO BRING AND DO PRIOR TO PROCEDURE:


- Please bring your PHOTO ID and INSURANCE CARD.
- Please bring a list of your current medications (including non-prescription) and allergies.
- **YOU MUST HAVE A DRIVER 18 YEARS OF AGE OR OLDER WHO WILL ACCEPT RESPONSIBILITY AND DRIVE YOU HOME.** They will be notified when your procedure is over and MUST be present at the time of discharge to receive instructions and teachings. You are NOT allowed to take a bus, taxi, walk home etc.
- If you do not have a ride, your procedure will be cancelled or rescheduled.
- You should plan to be at our facility for at least 2 hours.
- If you use any inhalers, please bring them with you to your appointment.
- Please read prep instructions THOROUGHLY.
- **DO NOT DRINK ANYTHING FOR 3 HOURS BEFORE YOUR PROCEDURE.**
- If your procedure is scheduled for the following times, then your driver must stay at the Wyoming Endoscopy Center. They may not leave the premises. Monday - Thursday at 4:00pm or 4:30pm Friday at 10:30am or 11:00am.

The Wyoming Endoscopy Center is located just North of Digestive Health Associates. The address is 7220 Commons Circle, Cheyenne WY 82009 and phone number is 307-462-0698.

Peg Solution Prep

1 Week Prior to your procedure:

- **PLEASE HOLD ANY GLP1 agonist for weight loss or diabetes for 1 week (tablets 24 hours) prior to your scheduled procedure.** These must be stopped prior to procedure due to increased risk for delayed gastric emptying. These medications include but are not limited to: Dulaglutide (TRULICITY), Exenatide (BYETTA), Exenatide Extended Release, (BYDUREON BCISE), Liraglutide, Saxenda (VICTOZA), Lixisenatide (ADLYXIN), Semaglutide, Wegovy, Rybelsus (OZEMPIC) and Tirzepatide (Mounjaro).
- If you take any of these medications for blood sugar control please contact your prescribing Provider for instructions. Not stopping these medications can result in cancellation of your procedure or require you to have a repeat procedure.



5 Days prior to your procedure: _____

- Discontinue Iron and Fish oil
- If on blood thinners please make us aware, but you will need to consult with prescribing physician for possible hold or alterations of blood thinner (Coumadin, Warfarin, Pradaxa, Xarelto and Eliquis)
- If you are **diabetic**, please see Diabetic Instructions
- Please pick up prescriptions from pharmacy (colon prep and optional anti-nausea medication called ondansetron) at your pharmacy. These prescriptions were sent electronically to your pharmacy at the time you were scheduled. If you do not pick these up within 7 days, your prescription may be filled and then put back on the shelf. Please call your pharmacy to verify the prescription is ready for pick up.
- You may take 4mg of Ondansetron every hour (up to 2 doses) with each prep dose for nausea.

3 Days prior to your procedure: _____


- Discontinue Fiber Supplements
- Begin a low-fiber Diet...please see Diets and Tips

1 Day prior to your procedure: _____

- Please have a light breakfast (soft egg, dried toast) before 9:00am, after 9:00am then you must begin a clear liquid diet...please see Diets and Tips
 - From 5-7 pm drink HALF the gallon of prep. Please pace yourself to drink about 8 oz every 15 minutes until the first half is completed.
 - Continue your clear liquid diet the rest of the night
 - Place the remaining prep in refrigerator until the next morning
- The Day of the procedure:

Beginning 5 Hours prior to your procedure _____, then please drink the second half of the gallon prep until the remaining prep is completely finished.

- You have a 2-hour time frame to complete the second half of the prep
- You **MUST** be finished with the entire prep 3 HOURS prior to your procedure.
- Please take your morning medications with a sip of water.



3 HOURS PRIOR TO YOUR PROCEDURE YOU MAY NOT HAVE ANYTHING BY MOUTH including Clear Liquids, Candy, Mints, Gum, no smoking or electronic cigarette. Please do not use chewing tobacco for 6 hours prior to your procedure. Failure to follow these instructions could result in your procedure being cancelled.

Please do not bring any valuable items to the Wyoming Endoscopy Center. We will not be held responsible for replacing any lost or misplaced items. Thank you.

DIET and TIPS

Bowel Prep is needed to perform an effective Procedure. If your bowel prep is not sufficient it will result in the need to reschedule the procedure as we will not be able to see lesions or polyps.

High Fiber foods to **AVOID** 3 days prior to procedure: _____

- whole wheat or whole grain breads, cereals, and pasta
- brown or wild rice, oats, kasha, barley, and quinoa
- Fruits (Dried and prune juice)
- seeds/nuts and foods containing them, peanut butter
- Dried beans, peas, and lentils • Coconut, Popcorn
- Vegetables (this includes both raw and undercooked)

Low Fiber ALLOWED:

- white bread
- white rice, plain white pasta, crackers
- refined hot cereals: such as cream of wheat or cold cereals less than 1 gram of fiber per serving
- pancakes/waffles made from refined flour
- fruit/vegetables juice with little or no pulp, fruit flavored drinks/flavored water
- tender meat, poultry, fish, eggs, and tofu
- milk and foods made from milk, yogurt, pudding, ice cream, cheese and sour cream-if tolerated
- butter, margarine, oils, and salad dressing without seeds



**Wyoming Endoscopy
Center**



**DIGESTIVE HEALTH
ASSOCIATES
of Cheyenne**

CLEAR LIQUID DIET

- water, sports drinks, energy drinks, clear sodas (7-up, sprite, ginger ale) tea, black coffee, juice without pulp
- Clear broths, gelatin, popsicles (yellow or green)
- sugar, honey, syrup, and hard candy
- PLEASE DO NOT EAT ANYTHING RED OR CONTAINING RED DYE
- avoid flavors such as: strawberry, cherry, watermelon, grape and orange
- **NO SOLID FOODS, NO MILK PRODUCTS, NO ORANGE JUICE and NO ALCOHOL**